

State and trends in the diversity, abundance and distribution of birds in Wellington City

May 2023



Authors:

NIKKI MCARTHUR, IAN FLUX AND ANNETTE HARVEY

State and trends in the diversity, abundance and distribution of birds in Wellington City.

Nikki McArthur¹, Ian Flux² and Annette Harvey³

¹ 17A Ida Street, Redwoodtown, Blenheim 7201

² PO Box 40694, Upper Hutt 5140

³ 35 Havelock Street, Mornington, Wellington 6021

This report was prepared for Greater Wellington Regional Council in fulfilment of the Contract for Services dated 12th November 2020.

30th May 2023.

Citation:

This report should be cited as:

McArthur, N.; Flux, I. and Harvey, A. 2023. State and trends in the diversity, abundance and distribution of birds in Wellington City. Client report prepared for Greater Wellington Regional Council, Wellington.

All photographs in this report are copyright © N. McArthur unless otherwise credited, in which case the person or organization credited is the copyright holder.

Cover Image: A tauhou / silvereye (*Zosterops lateralis*) perched on a karamū (*Coprosma robusta*) shrub in Zealandia Te Māra a Tāne Wildlife Sanctuary, 20th March 2017.

Image credit: Eric VanderWerf / Macaulay Library at the Cornell Lab of Ornithology ([ML53270701](https://www.macaulaylibrary.org/record/ML53270701)).

Table of Contents

Executive Summary.....	4
1. Introduction	6
2. Methods.....	9
2.1 Five-minute bird count data collection.....	9
2.2 Five-minute bird count data analysis.....	11
2.2.1 Trends in the encounter rates of native versus introduced species.....	11
2.2.2 Trends in the encounter rates of individual native forest bird species	12
2.2.3 Zealandia Te Māra a Tāne’s influence on native and introduced species richness.....	12
2.3 Citizen science data analysis	14
3. Results.....	16
3.1 Species diversity.....	16
3.2 Abundance and distribution of native forest bird species.....	18
3.2.1 Tūi (<i>Prosthemadera novaeseelandiae</i>)	20
3.2.2 Tauhou / Silvereye (<i>Zosterops lateralis</i>)	22
3.2.3 Riroriro / Grey warbler (<i>Gerygone igata</i>)	24
3.2.4 Pīwakawaka / New Zealand fantail (<i>Rhipidura fuliginosa</i>)	26
3.2.5 Kākā (<i>Nestor meridionalis</i>).....	28
3.2.6 Kererū (<i>Hemiphaga novaeseelandiae</i>).....	30
3.2.7 Pīpīwharau / Shining cuckoo (<i>Chrysococcyx lucidus</i>).....	32
3.2.8 Tīeke / North Island saddleback (<i>Philesturnus rufusater</i>).....	35
3.2.9 Kākāriki / Red-crowned parakeet (<i>Cyanoramphus novaezelandiae</i>).....	38
3.2.10 Korimako / Bellbird (<i>Anthornis melanura</i>).....	41
3.2.11 Pōpokotea / Whitehead (<i>Mohoua albicilla</i>)	44
3.2.12 Kōtare / New Zealand kingfisher (<i>Todiramphus sanctus</i>).....	47
3.2.13 Kārearea / New Zealand falcon (<i>Falco novaeseelandiae</i>).....	49
3.2.14 Toutouwai / North Island robin (<i>Petroica longipes</i>)	52
3.2.15 Tītipounamu / Rifleman (<i>Acanthisitta chloris</i>).....	55
3.2.16 Hīhi (<i>Notiomystis cincta</i>)	57
3.2.17 Miromiro / Tomtit (<i>Petroica macrocephala</i>)	59
3.2.18 Koekoeā / Long-tailed cuckoo (<i>Eudynamys taitensis</i>).....	61
3.2.19 Ruru / Morepork (<i>Ninox novaeseelandiae</i>).....	61
3.3 The Zealandia Te Māra a Tāne ‘halo effect’	64
4. Discussion.....	65

4.1	Changes in bird community composition, abundance and distribution.....	65
4.2	The role of citizen scientists in monitoring Wellington City’s bird fauna	67
5.	Recommendations	69
	Acknowledgements.....	70
	References	71
	Appendix One.....	75
	Appendix Two	81

Executive Summary

Five-minute bird counts have been carried out at 100 bird count stations in forest habitat throughout Wellington City's parks and reserves network each year between 2011 and 2022. The aim of these surveys is to monitor trends in the diversity, abundance and distribution of native forest birds throughout Wellington City's parks and reserves, to assess how local conservation management activities are impacting the city's native bird fauna.

Since 2011, the average number of native bird species being counted per bird count each year has risen by 41%, and the average number of introduced bird species being counted per count has increased by 47%. Between 2011 and 2022, average annual counts of kākā have increased by 260%, average annual counts of kererū have increased by 200%, average annual counts of tūī have increased by 85% and average annual counts of pīwakawaka / New Zealand fantails have increased by 49%. Over the same time period, average annual counts of tauhou / silvereyes have declined by 9%.

The predator-free Zealandia Te Māra a Tāne sanctuary is having a measurable 'halo' effect on native forest bird communities throughout Wellington City. The average number of native bird species recorded per five-minute bird count station declines with increasing distance from Zealandia Te Māra a Tāne's predator-proof fence. An average of 6.8 native bird species has been recorded at count stations closest to Zealandia Te Māra a Tāne's boundary between 2011 and 2022, whereas an average of only 4.6 native bird species were recorded at those count stations situated furthest from Zealandia Te Māra a Tāne's boundary over the same period. In contrast, the average number of introduced species recorded per five-minute bird count station increases with increasing distance from Zealandia Te Māra a Tāne's predator-proof fence. An average of 2.1 introduced bird species has been recorded at count stations closest to Zealandia Te Māra a Tāne between 2011 and 2022, whereas an average of 3.5 introduced bird species was recorded at count stations situated furthest away from the sanctuary over the same period.

These results suggest that the presence of large 'source' populations of native forest birds in Zealandia Te Māra a Tāne, together with the increasing extent and intensity of mammalian predator control being carried out in Wellington City, is driving spectacular recoveries in several previously rare or locally extinct native forest bird species in Wellington City. As a consequence, bird communities in native forest habitats in Wellington City parks and reserves are becoming more diverse, creating improved opportunities for local residents and visitors alike to encounter a wider range of New Zealand's native forest bird species in the heart of New Zealand's capital city. One clear sign of this increasing level of engagement is the large and growing number of citizen science bird records that are being collected in Wellington City, with a total of 95,761 observations of native forest birds having been reported to date by local residents and visitors since 2011.

It is recommended that Wellington City Council continues to carry out these five-minute bird counts on an annual basis, to monitor future improvements in the city's native bird communities as Wellington continues on its journey towards becoming the world's first predator-free capital city.

Keywords: Citizen science, eBird, encounter rate, five-minute bird count, generalised linear mixed-effect models, iNaturalist New Zealand, New Zealand Bird Atlas, Predator Free Wellington, Wellington City, Zealandia Te Māra a Tāne

Wellington City forest reserves bird health check

Low Concern

Large, stable or increasing populations.
Low to moderate predator risk.

Tūī, Tauhou / Silvereye, Riroriro / Grey warbler, Pīwakawaka / NZ fantail, Kākā, Kererū, Kōtare / NZ kingfisher



Moderate Concern

Small, localised or sparse populations.
Moderate to high predator risk.

Pīpīwharau / Shining cuckoo, Tīeke / North Island saddleback, Kākāriki / Red-crowned parakeet, Pōpokotea / Whitehead, Korimako / Bellbird, Kārearea / New Zealand falcon, Toutouwai / North Island robin, Hihi, Tītītipounamu / Rifleman



High Concern

Tiny or declining populations. Moderate to high predator risk.

Data deficient

Population size and trends poorly known.



Ruru / Morepork

Image credits: NZ Birds Online
(<http://nzbirdsonline.org.nz/>)

1. Introduction

Over the past twenty years there has been a dramatic improvement in the diversity, abundance and distribution of native forest birds in Wellington City (Miskelly *et al.* 2005; McArthur *et al.* 2021). This improvement is due to two key developments in the management of indigenous forest habitats in and around the city that have occurred in recent decades. Firstly, a series of species reintroductions to local predator-free sites such as Zealandia Te Māra a Tāne, Matiu/Somes Island and Mana Island have successfully established healthy source populations from which previously locally extinct or near-extinct bird species have been dispersing into nearby forested reserves in the city (Miskelly & Powlesland, 2013; McArthur *et al.* 2017). These species include kākā (*Nestor meridionalis*), kākārīki / red-crowned parakeet (*Cyanoramphus novaeseelandiae*), pōpokotea / whitehead (*Mohoua albicilla*) and korimako / bellbird (*Anthornis melanura*) (Miskelly *et al.* 2005; Froude, 2009; McLaughlin & Harvey, 2013; Miskelly, 2018). Secondly, ongoing multi-species mammalian predator control being carried out by Wellington City Council, Greater Wellington Regional Council and community conservation groups in many Wellington City parks and reserves and in surrounding suburban areas has resulted in local increases in resident native bird species such as tūī (*Prosthemadera novaeseelandiae*) and kererū (*Hemiphaga novaeseelandiae*) (Bell, 2008; Froude, 2009; Brockie & Duncan, 2012; McArthur *et al.* 2021) and is creating an opportunity for some of these recently reintroduced species to establish functional populations away from their original reintroduction sites.

Within Zealandia Te Māra a Tāne, a fenced, 225 ha predator-free sanctuary situated in the suburb of Karori, eleven endemic forest bird species have been reintroduced since the eradication of mammalian predators in 2000, and a further two species have recolonised of their own accord (Miskelly, 2018¹). This has led to the re-establishment of a diverse and abundant endemic forest bird community within Zealandia Te Māra a Tāne's predator-proof fence, which in turn has led to substantial declines in the abundance of three of the four native forest bird species that had been resident in Zealandia Te Māra a Tāne prior to the eradication of mammalian predators, namely tauhou / silvereye (*Zosterops lateralis*), riroriro / grey warbler (*Gerygone igata*) and pīwakawaka / fantail (*Rhipidura fuliginosa*) (Miskelly, 2018). At least six introduced bird species have also experienced similar, substantial declines within Zealandia Te Māra a Tāne over this time period, including chaffinch / pahirini (*Fringilla coelebs*), dunnoek (*Prunella modularis*) and song thrush / manu-kai-hua-rakau (*Turdus philomelos*) (Miskelly, 2018). These changes to Zealandia Te Māra a Tāne's bird community over the past two decades may foreshadow the changes that may occur in forest habitats in other Wellington City parks and reserves, as efforts to control and/or eradicate mammalian predators continue to expand and intensify.

Mammalian predator control and eradication efforts in Wellington City are continuing to grow in both intensity and coverage. Dozens of community groups across Wellington City are now participating in predator control activities, and in recent years the concept of creating 'predator free suburbs' has emerged. Following central government's announcement in July 2016 to work towards the goal of a Predator Free New Zealand by 2050, these Wellington City predator control efforts have further crystallised into the Predator Free Wellington initiative. Predator Free Wellington is being co-funded

¹ Miskelly (2018) lists 10 forest bird species reintroduced to Zealandia. In March 2019, an 11th species, tītīpounamu / rifleman (*Acanthisitta chloris*) was also reintroduced to the sanctuary (<https://www.facebook.com/ZEALANDIA/posts/10156377249456401>; accessed 04/04/2021).

by Wellington City Council, Greater Wellington Regional Council and the NEXT Foundation and plans to build on the proliferation of pest-free suburb projects with the aim of eradicating rats (*Rattus* spp.), mustelids (*Mustela* spp.) and possums (*Trichosurus vulpecula*) from Wellington City. The first stage of this project commenced in July 2019 with an effort to eradicate rats and mustelids from Miramar Peninsula, and preparations are now underway to begin an eradication of rats, possums and mustelids from an additional 14 suburbs from Island Bay to the CBD (<https://www.pfw.org.nz/>; accessed 04/04/2023). In 2018, the Capital Kiwi Project also began installing a network of predator traps on the Wellington Peninsula to the west of Wellington City, with the aim of creating a large area of predator-free habitat to allow North Island brown kiwi (*Apteryx mantelli*) to be re-established in the wild within Wellington City limits. By 2022, this trapping network had grown to over 4,500 traps across 23,500 hectares of land stretching from the Wellington south coast to Porirua. In November 2022 a major milestone was reached with the release of eleven North Island brown kiwi on Terawhiti Station, followed by the release of an additional 50 kiwi in May 2023 (<https://www.capitalkiwi.co.nz/>; accessed 25/05/2023).

These combined efforts to eradicate or control mammalian predators in Wellington City are likely to result in further dramatic changes in the distribution and abundance of native birds in Wellington. Monitoring the ongoing changes that are occurring to native bird populations in the city therefore provides a useful means by which the outcome of the considerable time and effort now being spent on improving Wellington City's biodiversity can be measured. For this reason, Wellington City Council identified a need to monitor local bird populations to provide a measure of the success or otherwise of their recently adopted Biodiversity Strategy & Action Plan (WCC, 2015). Goal 4.2.2a of this Biodiversity Strategy involves setting up a "consistent terrestrial outcome monitoring framework...incorporating existing monitoring work in a collaborative approach with other key organisations" (WCC, 2015).

Five-minute bird count monitoring was carried out between 2001 and 2009 in nine selected parks and reserves in Wellington City by Pacific Eco-Logic Ltd (Froude, 2009). These counts were successful in detecting substantial increases in the local abundance of tūi at a key time during which a major expansion in pest control efforts in Wellington City was underway. These counts also provided some of the earliest evidence that bird species re-introduced to Zealandia Te Māra a Tāne were dispersing and settling in nearby reserves (Froude, 2009). In 2011 this bird monitoring programme was replaced with a new survey designed to monitor changes in the distribution and abundance of native forest birds in forest habitats across the entire network of Wellington City parks and reserves, rather than only a selected subset of reserves (McArthur *et al.* 2012).

These counts have now been carried out each year since 2011 and have demonstrated the important influence that Zealandia Te Māra a Tāne has had on the native forest bird community in the wider Wellington City. Approximately a third of the native forest bird species being detected in Wellington City parks and reserves each year are species that have been re-introduced to Zealandia Te Māra a Tāne and have subsequently expanded their range to include a number of other parks and reserves in the city (McArthur *et al.* 2021). Furthermore, the ecological effects of the reintroduction of these species to Zealandia Te Māra a Tāne reported by Miskelly (2018) (namely, reductions in the abundance of several common, resident native and introduced bird species) have now been detected in forest habitats beyond Zealandia Te Māra a Tāne's predator-proof fence (McArthur *et al.* 2021). Individually, the bird species that have been reintroduced to Zealandia Te Māra a Tāne have had varying levels of success at colonising forest habitats elsewhere in the city. Both kākā and kākārīki / red-crowned parakeets are now regularly encountered in parks, reserves and suburban gardens throughout central Wellington, whereas tieke / North Island saddleback (*Philesturnus rufusater*), toutouwai / North Island

robin (*Petroica longipes*) and pōpokotea / whitehead all continue to have very sparse and localised distributions largely restricted to forest reserves immediately adjacent to Zealandia Te Māra a Tāne. These patterns indicate that mammalian predators are to varying degrees continuing to limit the ability of these species to establish self-sustaining populations in native forest habitats in the city (McArthur *et al.* 2021; Irwin & Empson, 2022). These counts have also shown that mean annual counts of native forest birds have increased by 41% between 2011 and 2022. This was driven largely by a 270% increase in mean annual counts of kākā, a 243% increase in mean annual counts of kererū, a 74% increase in mean annual counts of tūi and a 37% increase in mean annual counts of pīwakawaka / New Zealand fantails. This suggests that the ongoing steady expansion and intensification of mammalian predator control efforts throughout Wellington City since 2011 have led to measurable improvements in the population health of these particular bird species (McArthur *et al.* 2022).

Making use of the bird observations of local citizen scientists has allowed us to map the distribution of native birds in Wellington City in unprecedented detail. These maps have helped document the range expansion of recently re-introduced species such as kākā and kākārīki / red-crowned parakeet in Wellington City virtually in real time and have documented a number of local re-colonisation events that have occurred in recent years in several individual parks and reserves (McArthur *et al.* 2022).

This report provides an update on the emerging trends in the diversity, abundance and distribution of birds in forest habitats in parks and reserves throughout Wellington City, by analysing and reporting a twelfth year of five-minute bird count data, together with additional citizen-science data collected since the publication of the previous bird monitoring report in April 2022.

2. Methods

2.1 Five-minute bird count data collection

One hundred bird count stations were established at randomly selected locations in forest habitat in Wellington City parks and reserves in November 2011 and have been surveyed annually between 2011 and 2022 (Figure 2.1). These bird count stations were established at a minimum distance of 200 metres from one another and each station has been marked with either a blue triangle affixed to a living tree, or with pink flagging tape if situated in plantation forest.

Two five-minute bird counts have been carried out at each station each year, with the two counts at each station being carried out on a different day. All counts were carried out between early November and early January each year and counts were made only on fine, calm days between 1.5 hours after sunrise and 1.5 hours before sunset (approximately 7.30 am to 6.30 pm). At each station, an observer spent five minutes recording the number of individuals of all species seen or heard from the count station (i.e., an unbounded count as per Dawson & Bull, 1975 and Hartley & Greene, 2012). Care was taken not to record the same bird twice during a count. Two experienced observers conducted the counts each year, with each observer surveying approximately half of the bird count stations.

Bird conspicuousness can vary in response to external variables such as time of year, weather, time of day and change in observer (Bibby *et al.* 2000). To account for this, every effort was made to standardise or sample the range of variation in each of these factors to ensure that as much as possible any changes in the mean number of birds counted per station from one year to the next would be more likely to reflect changes in bird abundance rather than conspicuousness. Precautions taken include carrying out these counts during the same months each year and in similar weather conditions. Counts were carried out throughout the day, so sampled any variation in bird conspicuousness that occurred during the day.

Variation in the methods and abilities of observers can have a substantial impact on five-minute bird count results and can sometimes either mask or be mistaken for true changes in bird abundance or conspicuousness from one survey to the next (McArthur *et al.* 2013). For this reason, we've endeavoured to minimise the number of observers used to collect this five-minute bird count data, with only two changes being made so far during the twelve-year duration of this project. In each case, when one observer has been replaced with another, the second observer has remained the same across both years, thus providing us with some ability to differentiate observer-related variation in bird encounter rates from those caused by true changes in bird conspicuousness or abundance from one year to the next.




Figure 2.1: Locations of five-minute bird count stations surveyed annually in Wellington City parks and reserves between 2011 and 2022.

2.2 Five-minute bird count data analysis

The Wellington City five-minute bird count data were first entered into a Microsoft Excel spreadsheet and then imported into the statistical package R (R Core Team, 2021) for statistical analysis. Three separate analyses were carried out, the first of which looked at overall trends in the encounter rates of native and introduced birds over time; the second of which investigated temporal trends in the encounter rates of any individual species that occurred in at least 2.5% of counts (i.e., there were at least 50 individual detections during counts carried out between 2011 and 2022); and the third of which investigated the influence that Zealandia Te Māra a Tāne's proximity has on the diversity of native and introduced bird species recorded at a bird count station.

For all three analyses, generalised linear mixed-effects models with a Poisson error were used to investigate temporal and spatial trends in bird encounter rates. For each analysis, we compared models from a candidate set to determine which of the plausible relationships between the number of birds encountered and time best explained the data. The model that fitted the data best was selected using Akaike's Information Criterion corrected for small sample size (AICc; Burnham and Anderson, 2002), with the model with the lowest AICc value providing the best fit to the data using the lowest number of model parameters. We then used our preferred models to assess whether trends in bird encounter rates were increasing or decreasing over time by calculating estimates of slope, and 95% confidence intervals (CIs) around estimates. Positive estimates of slope indicated increases in bird encounter rates, while negative estimates of slope indicated decreases in bird encounter rates. We used the 'lme4' package (Bates *et al.* 2015) in the statistical package R (R Core Team, 2021) for all three analyses.

2.2.1 Trends in the encounter rates of native versus introduced species

To analyse overall trends in the encounter rates of native versus introduced birds over time, three plausible models were considered:

1. **No change:** Bird encounter rates have not changed between 2011 and 2022 but encounter rates of native and introduced species have differed. This model included the number of individuals of all species detected during a count in a given year as a response variable, and the status of each species (native or introduced) as a predictor variable.
2. **Same population trend for both native and introduced species:** Bird encounter rates have changed between 2011 and 2022, at the same rate for both native and introduced species. This model consisted of the same response variable as the "no change" model above but included both species status and year as predictor variables.
3. **Different population trends for native versus introduced species:** Bird encounter rates have changed between 2011 and 2022, but at different rates for native versus introduced species. This model consisted of the same response and predictor variables as the "same population trend" model above but included an additional interaction term between the status and year predictor variables to allow for the slope of the relationship with time to vary between native and introduced species.

For each of these three models, a random intercept term was included for each station to account for the repeated-measures design. We did not include observer in the model because observer was partly confounded with station. However, only four observers have been used over the eleven-year series of counts, and generally at least five levels of a random effect (in this case, the observer) are required to achieve robust estimates of variance (Gelman & Hill, 2007).

2.2.2 Trends in the encounter rates of individual native forest bird species

To analyse trends in the encounter rates of individual bird species between 2011 and 2022, two plausible models were considered:

1. **No trend:** Bird encounter rates have not changed between 2011 and 2022.
2. **Trend over time:** Bird encounter rates have either increased or decreased between 2011 and 2022.

Both models included the number of individuals of each species detected during a count in a given year as a response variable, an intercept term and a random intercept term for each station to account for the repeated-measures design. The “trend over time” model also included year as a predictor variable. As with the first analysis, observer was not included as a random effect. Where it was necessary, an observation-level random effect was also included to account for overdispersion, i.e., a higher error variance than assumed by the Poisson error distribution for these models (Harrison, 2014).

2.2.3 Zealandia Te Māra a Tāne’s influence on native and introduced species richness

To analyse how proximity to Zealandia Te Māra a Tāne influences the number of native or introduced species recorded at a bird count station, three plausible models were considered:

1. **No spatial trend:** Differences in species richness occur between native and introduced species at any given location but show no relationship with distance from Zealandia Te Māra a Tāne.
2. **Same spatial trend for native and introduced species:** Differences in species richness occur between native and introduced species at any given location, and these differences vary in relation to the distance of the count station from Zealandia Te Māra a Tāne at the same rate for both native and introduced species.
3. **Different spatial trends for native versus introduced species:** Differences in species richness occur between native and introduced species at any given location, and these differences vary in relation to the distance of the count station from Zealandia Te Māra a Tāne but at different rates for native and introduced species.

All three models contained species richness (i.e., the mean number of species detected at each station) as a response variable, an intercept term, and a random intercept term for each station (to account for the repeated-measures design) and for year (to account for variation between years). All models also included a log-transformed offset term consisting of the number of native and introduced

species included in the analysis, to account for there being more native than introduced species in the analysis. A linear relationship between species richness and distance to Zealandia Te Māra a Tāne was also assumed. Model one (no spatial trend) included status (whether a species is native or introduced) as a predictor variable. Model two (same spatial trend for native and introduced species) included status and distance to the nearest point of Zealandia Te Māra a Tāne's boundary as predictor variables. Model three (different spatial trends for native and introduced species) included the same predictor variables as model two, but also included an interaction term between status and distance to Zealandia Te Māra a Tāne as an additional predictor variable to allow for the spatial trend in species richness to vary between native and introduced species.

Patterns in the distribution of native birds among Wellington City reserves were also examined by mapping the relative frequency at which each native forest bird species was detected at each bird count station using ArcMap version 10.8.2. Although this technique does not explicitly take into account relative differences in abundance (e.g., less common species present within sight or earshot of a bird count station are less likely to be detected) or variation in detection probabilities between species (e.g., less conspicuous species will also be less likely to be detected), it should be sufficient to detect relatively conspicuous patterns in species' distributions and habitat use (Mackenzie *et al.* 2006).

2.3 Citizen science data analysis

As a result of the increasing popularity of citizen science, there is a rapidly growing pool of bird observation data collected by citizen scientists available online which can be combined with our more systematic five-minute bird count data to help detect changes in bird distribution in Wellington City over time. To date, residents and visitors to Wellington City have contributed over 270,000 bird observations to online databases and citizen science projects such as the New Zealand eBird database, the New Zealand Bird Atlas, iNaturalist and the Great Kererū Count.

The New Zealand eBird database (<http://ebird.org/content/newzealand/>), which in turn hosts the New Zealand Bird Atlas dataset, is run by the Cornell Lab of Ornithology in partnership with Birds New Zealand / Te Kāhui Mātai Manu o Aotearoa (formerly the Ornithological Society of New Zealand). It provides a facility for recreational birdwatchers and professional ornithologists to permanently record their bird observations in a standard format in one centralised location and makes these observations available to researchers, conservation managers and environmental policy makers (Scofield *et al.* 2012). Globally, the eBird database is now the largest and fastest growing biodiversity database in the world, with over 820,000 unique users having so far contributed over 1.3 billion bird records describing the distribution of 98% of the world's bird species (Sullivan *et al.* 2014; <https://ebird.org/news/2022-year-in-review>, accessed 29/04/2023).

Within the eBird database, automated data filters and an expert review process ensure that these data are of high quality and accuracy (Sullivan *et al.* 2014). We used eBird's "download data" tool to access the March 2023 release of the eBird Basic Dataset (EBD) and to build custom datasets containing citizen science records of all native forest bird species recorded in Wellington City between 2011 and 2023. We formatted these datasets using Microsoft Excel, including removing any extraneous data fields and converting latitude/longitude coordinates to New Zealand Transverse Mercator (NZTM2000) coordinates. We then saved these files as .csv files so that they could be imported into ArcMap and converted into shapefiles. Once in ArcMap, we visually inspected these eBird records to locate and remove any records containing obvious location errors (e.g., records placed offshore, or for which location descriptions didn't match the coordinates provided) or absence records, before adding these records to the distribution maps created from the five-minute bird count data. A total of 89,104 records of native forest birds observed in Wellington City was retrieved from eBird using this process, representing 93% of all of the citizen science bird observations included in this report.

The iNaturalist New Zealand database is the second-largest online source of citizen science bird data for the Wellington region. iNaturalist New Zealand is a database that allows citizen scientists to submit, share and store natural history observations online, and unlike eBird it is designed to accept records for almost any taxon of plant or animal rather than just birds. iNaturalist New Zealand (<https://inaturalist.nz/>) is run by a charitable trust called the New Zealand Bio-recording Network Trust and was established using funding from the New Zealand Government's Terrestrial Freshwater Biodiversity Information System Fund.

Within the iNaturalist New Zealand database, a community peer-review process is used to validate records, with records tagged as either "research grade" or "casual grade" depending on whether or not the original species identifications have been verified by other iNaturalist New Zealand users. We used the search tool on the iNaturalist New Zealand website (<https://inaturalist.nz/>) to create and download a custom dataset of all bird observations recorded in Wellington City between 2011 and 2022. We formatted this dataset using Microsoft Excel, then saved the resulting file as a .csv file so

that it could be imported into ArcMap and converted to a shapefile. We then displayed the data on a map and visually inspected them, removing records with obvious location errors, before adding these records to the distribution maps created from the five-minute bird count data. A total of 6,083 records of native forest birds observed in Wellington City was retrieved from iNaturalist New Zealand using this process, representing 6% of all of the citizen science bird observations included in this report.

Kererū Discovery's Great Kererū Count dataset is the third-largest source of citizen science bird data available for Wellington. The Great Kererū Count was a nationwide kererū survey that took place each year between 2014 and 2021 (Hartley 2021). Observers from around the country were encouraged to record the presence or absence of kererū at locations of their choosing over a 10-day period each September. In 2021, 12,002 reports were received nationwide, with a total of 24,562 kererū counted (<https://kererudiscovery.org.nz/the-great-kereru-count/>; accessed 21/04/2022). We made a request for access to the Great Kererū Count data from Kererū Discovery and received a .csv file containing 574 observations reported from Wellington City. We imported this .csv file into ArcMap and visually inspected the records to locate and remove any records containing obvious location errors before adding these records to the distribution maps created from the five-minute bird count data. The 574 kererū observations sourced from the Great Kererū Count dataset represent 1% of all of the citizen science bird observations included in this report.

A key difference between these citizen science datasets and the five-minute bird count data is that the temporal and spatial distribution of search effort spent by citizen scientists varies unpredictably from year to year, whereas this search effort is standardised during these five-minute bird counts. Nonetheless, accurate bird observations submitted by citizen scientists have the potential to complement distribution data derived from our five-minute bird count dataset by providing information describing the presence of native forest birds at locations and in habitats not sampled by these five-minute bird counts. By collecting bird observation data all year round, local citizen scientists are also better placed to record local colonisation events prior to the colonising species being detected during these five-minute bird counts.

3. Results

3.1 Species diversity

Fifty bird species have been detected during five-minute bird counts carried out in Wellington City parks and reserves between 2011 and 2022. This includes two species (tītīpounamu / rifleman and tākupu / Australasian gannet; *Morus serrator*) that were detected for the first time in 2022. A full list of the 50 bird species detected, along with their current national and regional conservation status rankings, can be found in Appendix One of this report.

Of these 50 bird species, 24 are typically found in native forest habitats, including 17 native or endemic species and seven introduced and naturalised species. The remaining 26 bird species recorded during these counts are marine, coastal or open country species such as kororā / little penguin (*Eudyptula minor*), tōrea pango / variable oystercatcher (*Haematopus unicolor*) and kāhu / swamp harrier (*Circus approximans*) and are not included in any of the individual species' analyses reported below. Among the 17 native or endemic bird species recorded during these counts, one is ranked as 'Threatened - Nationally Vulnerable', one is ranked as 'Threatened - Nationally Increasing', three are ranked as 'At Risk - Nationally Declining', one is ranked as 'At Risk – Nationally Relict', two are ranked as 'At Risk - Nationally Recovering', and nine are ranked as 'Not Threatened' under the New Zealand Threat Classification Scheme (Appendix One; Robertson *et al.* 2021).

The best model to describe the overall trends in encounter rates of native versus introduced birds in native forest habitat in Wellington City parks and reserves over time was the model that assumed that population trends would differ between native versus introduced species. This model had a lower AICc value compared to the next closest model considered ($\Delta AICc = 6.97$). According to this model, the mean number of native birds encountered per count has increased between 2011 and 2022 (the coefficient of year on log abundance was 0.22; 95% CI 0.20 - 0.25). Forty-one percent more native birds were encountered in 2022 than in 2011, with a mean of 6.9 (± 0.2 SE) native birds encountered per bird count in 2021 compared to 4.9 (± 0.2 SE) native birds per count in 2011. The mean number of introduced birds encountered per count has also increased between 2011 and 2022 (the coefficient of year on log abundance was 0.15; 95% CI 0.10–0.20). Forty-seven percent more introduced birds were counted in 2021 than in 2011, with a mean of 5.6 (± 0.3 SE) introduced birds encountered per bird count in 2021 compared to 3.8 (± 0.1 SE) introduced birds per count in 2011 (Figure 3.1).




Figure 3.1: Trends in the mean number of native and introduced birds encountered per bird count in Wellington City parks and reserves between 2011 and 2022. Individual data points (\pm standard error) represent the mean number of native and introduced bird species detected per count each year. Solid orange and dashed blue lines (\pm 95% Confidence Intervals) represent the modelled trend in the number of native and introduced bird species recorded per count each year.

3.2 Abundance and distribution of native forest bird species

Of the 19 native forest bird species that have been recorded in Wellington City since 2011 (comprising 17 recorded during these five-minute bird counts and a further two species reported by citizen scientists), 11 species were encountered on at least 50 occasions during these five-minute bird counts (2.5% of all counts). Based on the lowest AICc values of the two candidate models, seven of these 11 species showed trends over time, whereas four species showed no trend. For species where year has an effect on mean encounter rates, a coefficient estimate >0 indicates an increase in encounter rates over time, with greater coefficient estimates indicating faster rates of increase. Conversely, a negative coefficient estimate indicates a decrease in encounter rates over time. Based on the coefficient estimates for the seven native forest bird species that showed trends over time, five of these species (tūī, pīwakawaka / New Zealand fantail, kākā, kererū and tīeke / North Island saddleback) showed an increase in mean encounter rates over time, and two species (tauhou / silvereye and pīpīwharau / shining cuckoo; *Chrysococcyx lucidus*) showed a decrease in encounter rates (Table 3.1).

The following individual species accounts are listed in decreasing order of their current or recent abundance in Wellington City. Species that are most frequently encountered during the five-minute bird counts are covered first, and the species that are only seldom encountered, or not encountered at all during these five-minute bird counts are treated last. Every species of native forest bird that has been observed in Wellington City outside of Zealandia since 2011 is included in this section of the report. A separate summary table of annual native forest bird encounter rates can be found in Appendix Two of this report.

Table 3.1: Summary of models of population change in 11 native forest bird species detected on at least 50 occasions in Wellington City parks and reserves since 2011, and trend if detected. The ‘selected model’ column indicates which model was selected for each species. The ‘trend’ column indicates whether the change in abundance was increasing, decreasing, or if no trend was detected (‘-’) based on estimates of slope.

Species	AICc model 1	AICc model 2	Δ AICc	Selected model	Trend	Coefficient estimate (95% CIs)
Tūi	8526.79	8347.67	179.12	Model 2 (Trend over time)	Increasing	0.060 (0.051 – 0.069)
Pīwakawaka NZ fantail	3537.30	3497.73	39.57	Model 2 (Trend over time)	Increasing	0.068 (0.046 – 0.086)
Kākā	2193.00	2103.75	89.25	Model 2 (Trend over time)	Increasing	0.144 (0.112 – 0.169)
Kererū	1758.82	1709.09	49.73	Model 2 (Trend over time)	Increasing	0.131 (0.096 – 0.167)
Tieke North Island saddleback	732.52	731.52	1.00	Model 2 (Trend over time)	Increasing?	0.045 (-0.008 – 0.092)
Riroriro Grey warbler	6408.44	6409.49	1.05	Model 1 (No change over time)	-	-
Pōpokotea Whitehead	709.70	711.64	1.94	Model 1 (No change over time)	-	-
Kōtare New Zealand kingfisher	609.78	611.67	1.89	Model 1 (No change over time)	-	-
Kākāriki Red-crowned parakeet	697.89	697.94	0.04	Model 1 (No change over time)	-	-
Tauhou Silvereye	8743.35	8733.58	9.77	Model 2 (Trend over time)	Decreasing	-0.018 (-0.029 – -0.008)
Pīpīwharauoa Shining cuckoo	2293.22	2283.49	9.73	Model 2 (Trend over time)	Decreasing	-0.055 (-0.090 – -0.024)

3.2.1 Tūī (*Prosthemadera novaeseelandiae*)

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The tūī is the native forest bird species that is most frequently encountered during these five-minute bird counts. Tūī encounter rates have increased by 85% since 2011, from a mean of 1.35 tūī recorded per five-minute bird count in 2011 to a mean of 2.50 tūī per count in 2022 (the coefficient of year on log abundance was 0.060; 95% CI 0.051 – 0.069; Figure 3.2).



Image courtesy of Tony Whitehead/NZ Birds Online

The tūī is the native forest bird species most commonly reported by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 23,145 encounters with tūī, demonstrating that this species is now common and widespread in native forest, suburban and urban habitats throughout the city (Figure 3.3).




Figure 3.2: Mean (±SE) number of tūī recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.3: Distribution of tūī in Wellington City between 2011 and 2023. Orange circles represent tūī detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of tūī detected per station between 2011 and 2022. Yellow and green circles represent tūī observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.2 Tauhou / Silvereye (*Zosterops lateralis*)

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The tauhou / silvereye is the second most frequently encountered native forest bird species in these five-minute bird counts. Tauhou / silvereye encounter rates have declined by 9% since 2011, from a mean of 1.86 birds recorded per five-minute bird count in 2011 to 1.69 birds recorded per count in 2022 (the coefficient of year on log abundance was -0.018; 95% CI -0.029 – -0.008 Figure 3.4).



Image courtesy of Ormond Torr/NZ Birds Online

The tauhou / silvereye is the third most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 11,432 encounters with tauhou / silvereyes, demonstrating that this species is common and widespread in native forest, suburban and urban habitats throughout the city (Figure 3.5).




Figure 3.4: Mean (\pm SE) number of tauhou / silvereyes recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.5: Distribution of tauhou / silvereyes in Wellington City between 2011 and 2023. Orange circles represent tauhou / silvereye detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of tauhou / silvereyes detected per station between 2011 and 2022. Yellow and green circles represent tauhou / silvereye observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.3 Riroriro / Grey warbler (*Gerygone igata*)



Image courtesy of Bartek Wypych/NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The riroriro / grey warbler is the third most frequently encountered native forest bird species in these five-minute bird counts. Despite some moderate year-to-year fluctuations, riroriro / grey warbler encounter rates have not changed significantly between 2011 and 2022. Riroriro / grey warbler encounter rates have varied between a low of 0.84 riroriro / grey warblers per count in 2011 and a high of 1.45 per count in 2015 (Figure 3.6).

The riroriro / grey warbler is the fourth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 10,436 encounters with riroriro / grey warblers, demonstrating that this species is common and widespread in native forest and suburban habitats, but is largely absent from urban habitats throughout the city (Figure 3.7).




Figure 3.6: Mean (±SE) number of riroriro / grey warblers recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.7: Distribution of riroriro / grey warblers in Wellington City between 2011 and 2023. Orange circles represent riroriro / grey warbler detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of riroriro / grey warblers detected per station between 2011 and 2022. Yellow and green circles represent riroriro / grey warbler observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.4 Pīwakawaka / New Zealand fantail (*Rhipidura fuliginosa*)



Image courtesy of Cheryl Marriner/NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The pīwakawaka / New Zealand fantail is the fourth most frequently encountered native forest bird species in these five-minute bird counts. Pīwakawaka / New Zealand fantail encounter rates have increased by 49% since 2011, from a mean of 0.35 pīwakawaka / New Zealand fantails recorded per five-minute bird count in 2011 to a mean of 0.52 pīwakawaka / New Zealand fantails per count in 2022 (the coefficient of year on log abundance was 0.068; 95% CI 0.046 – 0.086; Figure 3.2.7). Moreover, pīwakawaka / New Zealand fantails

have increased by 148% over the past seven years, from a low of 0.21 bird per count in 2015, to 0.52 birds per count in 2022 (Figure 3.8).

The pīwakawaka / New Zealand fantail is the fifth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 8,564 encounters with pīwakawaka / New Zealand fantails, demonstrating that this species is common and widespread in native forest, suburban and urban habitats throughout the city (Figure 3.9).




Figure 3.8: Mean (\pm SE) number of pīwakawaka / New Zealand fantails recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.9: Distribution of pīwakawaka / New Zealand fantails in Wellington City between 2011 and 2023. Orange circles represent pīwakawaka / New Zealand fantail detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of pīwakawaka / New Zealand fantails detected per station between 2011 and 2022. Yellow and green circles represent pīwakawaka / New Zealand fantail observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.5 Kākā (*Nestor meridionalis*)

National conservation status:

At Risk, Recovering (Robertson *et al.* 2021).

Regional conservation status:

At Risk, Recovering (Crisp, 2020).

The kākā was reintroduced to Zealandia Te Māra a Tāne between 2002 and 2007 (Miskelly, 2018) and is now the fifth most frequently encountered native forest bird species in these five-minute bird counts. Kākā encounter rates have increased by 260% since 2011, from a mean of 0.10 kākā recorded per five-minute bird count in 2011 to a mean of 0.36 kākā per count in 2022 (the coefficient of year on log abundance was 0.144; 95% CI 0.112 – 0.169; Figure 3.10).



Image courtesy of Jean-Claude Stahl/NZ Birds Online

The kākā is now the second most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 12,851 encounters with kākā, demonstrating that this species is now common and widespread in native forest and suburban habitats in the central and southern suburbs of Khandallah, Ngaio, Wadestown, Karori, Kelburn, Te Aro, Mt Cook, Brooklyn, Newtown and Island Bay. Kākā are also continuing to spread gradually northwards into Johnsonville and Tawa, and eastwards onto Miramar Peninsula (Figure 3.11).




Figure 3.10: Mean (\pm SE) number of kākā recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.11: Distribution of kākā in Wellington City between 2011 and 2023. Orange circles represent kākā detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of kākā detected per station between 2011 and 2022. Yellow and green circles represent kākā observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.6 Kererū (*Hemiphaga novaeseelandiae*)

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: At Risk, Recovering (Crisp, 2020).

The kererū is the sixth most frequently encountered native forest bird species in these five-minute bird counts. Kererū encounter rates have increased by 200% since 2011, from a mean of 0.07 kererū recorded per five-minute bird count in 2011 to a mean of 0.21 kererū per count in 2022 (the coefficient of year on log abundance was 0.131; 95% CI 0.096 – 0.167; Figure 3.12). Significant ‘hotspots’ in kererū encounter rates occur in Ōtari-Wilton’s Bush and in Khandallah Park, two Wellington City reserves that contain areas of old-growth native forest (Figure 3.13).



Image courtesy of Arindam Bhattacharya/
NZ Birds Online

The kererū is the sixth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 7,174 encounters with kererū, demonstrating that kererū are widespread in native forest and suburban habitats throughout the western and central suburbs of the city and on northern Miramar Peninsula, but are less frequently reported in northern, southern and eastern suburbs (Figure 3.13).




Figure 3.12: Mean (\pm SE) number of kererū recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.13: Distribution of kererū in Wellington City between 2011 and 2023. Orange circles represent kererū detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of kererū detected per station between 2011 and 2022. Yellow, green and blue circles represent kererū observations reported by local citizen scientists via eBird, iNaturalist and the Great Kereru Count between 2011 and 2023.

3.2.7 Pīpīwharau / Shining cuckoo (*Chrysococcyx lucidus*)



Image courtesy of Rob Lynch/NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The pīpīwharau / shining cuckoo is the seventh most frequently encountered native forest bird species in these five-minute bird counts. Pīpīwharau / shining cuckoo encounter rates have declined in Wellington City between 2011 and 2022, with an annual mean of 0.21 pīpīwharau / shining cuckoos recorded per five-minute bird count between 2011 and 2018 compared to an annual mean of 0.11 pīpīwharau / shining cuckoos recorded per count between 2019 and 2022 (the coefficient of year on log abundance was

-0.055; 95% CI -0.090 - -0.024; Figure 3.14). A sudden and abrupt decline in pīpīwharau / shining cuckoo counts was observed between 2018 and 2019, followed by a slow recovery since 2019 (Figure 3.14).

Pīpīwharau / shining cuckoos are sparsely distributed throughout Wellington City, though 'hotspots' in encounter rates appear to occur in forested reserves adjacent to Zealandia, and in Khandallah Park (Figure 3.15). The pīpīwharau / shining cuckoo is the fifteenth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 1,247 encounters with pīpīwharau / shining cuckoos, demonstrating that these birds are sparsely distributed in native forest and occur more infrequently in suburban habitats throughout the city (Figure 3.15).




Figure 3.14: Mean (\pm SE) number of pīpīwhararua / shining cuckoos recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.15: Distribution of pīpīwharau / shining cuckoos in Wellington City between 2011 and 2023. Orange circles represent pīpīwharau / shining cuckoo detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of pīpīwharau / shining cuckoos detected per station between 2011 and 2022. Yellow and green circles represent pīpīwharau / shining cuckoo observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.8 Tīeke / North Island saddleback (*Philesturnus rufusater*)

National conservation status:

At Risk, Recovering (Robertson *et al.* 2021).

Regional conservation status:

Regionally Vulnerable (Crisp, 2020).

The tīeke / North Island saddleback was reintroduced to Zealandia Te Māra a Tāne between 2002 and 2003 (Miskelly, 2018) and is now the tenth most frequently encountered native forest bird species in these five-minute bird counts. Tīeke / North Island saddleback encounter rates have increased since 2011, from a low of 0.02 tīeke / North Island saddleback recorded per five-minute bird count in 2012 to a mean of 0.07 tīeke / North Island saddleback per count in 2022 (the coefficient of year on log abundance was 0.045; 95% CI - 0.008 - 0.092; Figure 3.16). Tīeke / North Island saddleback encounter rates have fluctuated moderately from year-to-year however, and the confidence limits around the coefficient estimate include a lower CI less than zero, so the certainty around this increase in encounter rates is currently low. Tīeke / North Island saddleback are largely restricted to Zealandia Te Māra a Tāne and to forested reserves less than 1-2 km from Zealandia Te Māra a Tāne’s pest-proof boundary fence, so this increase in encounter rates may be an indication that the intensification of mammalian predator control efforts in Polhill Reserve in recent years is benefitting this species.



Image courtesy of Rob Lynch/NZ Birds Online

The tīeke / North Island saddleback is the seventh most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 4,139 encounters with tīeke / North Island saddlebacks and the majority of these sightings were reported either from inside Zealandia Te Māra a Tāne or in native forest habitats within 1 km of Zealandia Te Māra a Tāne’s boundary fence (Figure 3.17). This localised distribution of tīeke / North Island saddleback observations strongly suggests that one or more environmental factors are severely limiting the ability of this species to colonise forest habitats beyond Zealandia Te Māra a Tāne’s predator-proof fence.




Figure 3.16: Mean (\pm SE) number of tīeke / North Island saddlebacks recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.17: Distribution of tīeke / North Island saddlebacks in Wellington City between 2011 and 2023. Orange circles represent tīeke / North Island saddleback detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of tīeke / North Island saddlebacks detected per station between 2011 and 2022. Yellow circles represent tīeke / North Island saddleback observations reported by local citizen scientists via eBird between 2011 and 2023.

3.2.9 Kākāriki / Red-crowned parakeet (*Cyanoramphus novaezelandiae*)



Image courtesy of Laurie Ross/NZ Birds Online

National conservation status: At Risk, Relict (Robertson *et al.* 2021).

Regional conservation status: At Risk, Recovering (Crisp, 2020).

The kākāriki / red-crowned parakeet was reintroduced to Zealandia Te Māra a Tāne between 2010 and 2011, although it is likely that this species had begun re-colonising Wellington City as early as 1999, following the eradication of rats from Kāpiti Island in 1996 and the introduction of kākāriki / red-crowned parakeets to Matiu/Somes Island in 2003 (Miskelly *et al.* 2005). Kākāriki / red-crowned parakeets are now the eleventh most frequently encountered native forest bird species in these five-minute bird counts. Kākāriki / red-crowned parakeet encounter rates have not changed significantly between

2011 and 2022, but have varied between a low of 0.01 birds counted per five-minute bird count in 2011, 2012 and 2013 and a high of 0.09 birds per count in 2015 and 2019 (Figure 3.18). Beyond Zealandia Te Māra a Tāne, kākāriki / red-crowned parakeets now appear to be either established or regular visitors to several forested reserves in the city, including in Wright’s Hill Reserve, Polhill Reserve, Ōtari-Wilton’s Bush, Khandallah Park, Huntleigh Park and the Wellington Botanic Gardens (Figure 3.19).

Kākāriki / red-crowned parakeets are the twelfth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 2,046 encounters with kākāriki / red-crowned parakeets demonstrating that this species is sparsely distributed in both native forest and suburban habitats in central and western Wellington suburbs and is now an occasional visitor to forest habitats on Miramar Peninsula (Figure 3.19).




Figure 3.18: Mean (\pm SE) number of kākāriki / red-crowned parakeets recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.19: Distribution of kākāriki / red-crowned parakeets in Wellington City between 2011 and 2023. Orange circles represent kākāriki / red-crowned parakeet detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of kākāriki / red-crowned parakeets detected per station between 2011 and 2022. Yellow and green circles represent kākāriki / red-crowned parakeet observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.10 Korimako / Bellbird (*Anthornis melanura*)



Image courtesy of Craig McKenzie/NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The korimako / bellbird was reintroduced to Zealandia Te Māra a Tāne between 2001 and 2011 (Miskelly 2018), although at least two individuals were present in the suburb of Karori during 1999 and 2000 (Miskelly *et al.* 2005). The korimako / bellbird is now the thirteenth most frequently encountered native forest bird species in these five-minute bird counts. Korimako / bellbirds have only been detected on 48 occasions during these five-minute bird counts, providing too few detections to allow trends in encounter rates to be modelled. Korimako

/ bellbird encounter rates have fluctuated from year to year between 2011 and 2022, from zero birds recorded in 2011 and 2015, to a mean of 0.06 birds encountered per bird count in 2022. Despite these year-to-year fluctuations, there does not appear to have been any overall trend in bellbird encounter rates between 2011 and 2022 (Figure 3.20).

The korimako / bellbird is the tenth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 2,196 encounters with korimako / bellbirds, demonstrating that bellbirds are very sparsely distributed in native forest and suburban habitats across Wellington City. A resident population is now established in Zealandia Te Māra a Tāne (Miskelly, 2018) and possibly also in the Wellington Botanic Gardens, Ōtari-Wilton's Bush and in Khandallah Park (Figure 3.21).




Figure 3.20: Mean (\pm SE) number of korimako / bellbirds recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.21: Distribution of korimako / bellbirds in Wellington City between 2011 and 2023. Orange circles represent korimako / bellbird detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of korimako / bellbirds detected per station between 2011 and 2022. Yellow and green circles represent korimako / bellbird observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.11 Pōpokotea / Whitehead (*Mohoua albicilla*)



Image courtesy of Tony Whitehead/NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The pōpokotea / whitehead was reintroduced to Zealandia Te Māra a Tāne between 2001 and 2002 (Miskelly, 2018) and is now the eighth most frequently encountered native forest bird species in these five-minute bird counts. Despite some minor year-to-year fluctuations, pōpokotea / whitehead encounter rates have not changed significantly between 2011 and 2022.

Pōpokotea / whitehead encounter rates have varied between a high of 0.09 pōpokotea / whiteheads per count in 2012 and a low of 0.04 per count in 2013 (Figure 3.22).

The pōpokotea / whitehead is the ninth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 3,360 encounters with pōpokotea / whiteheads. These records show that whiteheads are largely restricted to Zealandia Te Māra a Tāne and to forest reserves within 1-2 km of Zealandia Te Māra a Tāne's boundary fence, however they have been recorded further afield on occasions, including in Trelissick Park, Ōtari-Wilton's Bush, Tinakori Hill, Wellington Botanical Gardens, Mākara Peak and Prince of Wales Park (Figure 3.23). This localised distribution of pōpokotea / whitehead observations strongly suggests that one or more environmental factors are severely limiting the ability of this species to colonise forest habitats beyond Zealandia Te Māra a Tāne's predator-proof fence.




Figure 3.22: Mean (\pm SE) number of pōpokotea / whiteheads recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.23: Distribution of pōpokotea / whiteheads in Wellington City between 2011 and 2023. Orange circles represent pōpokotea / whitehead detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of pōpokotea / whiteheads detected per station between 2011 and 2022. Yellow and green circles represent pōpokotea / whitehead observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.



Image courtesy of Bartek Wypych/NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The kōtare / New Zealand kingfisher is the ninth most frequently encountered native forest bird species in these five-minute bird counts. Despite minor year-to-year fluctuations, kōtare / New Zealand kingfisher encounter rates have not changed significantly between 2011 and 2022. Kōtare / New Zealand kingfisher encounter rates have varied between a low of 0.02 kōtare / New Zealand kingfishers per count in 2016 and a high 0.06 per count in 2019 (Figure 3.24).

The kōtare / New Zealand kingfisher is the thirteenth most frequently reported native forest bird species by citizen scientists in Wellington City. Between 2011 and 2023, citizen scientists reported a total of 1,865 encounters with kōtare / New Zealand kingfishers, demonstrating that kingfishers are sparsely distributed in native forest, suburban and coastal habitats throughout the city, but with a particular concentration of records in the northern, eastern and southern parts of Miramar Peninsula (Figure 3.25).




Figure 3.24: Mean (\pm SE) number of kōtare / New Zealand kingfishers recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.25: Distribution of kōtare / New Zealand kingfishers in Wellington City between 2011 and 2023. Orange circles represent kōtare / New Zealand kingfisher detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of kōtare / New Zealand kingfishers detected per station between 2011 and 2022. Yellow and green circles represent kōtare / New Zealand kingfisher observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.13 Kārearea / New Zealand falcon

(*Falco novaeseelandiae*)



Image courtesy of Steve Attwood/NZ Birds Online

National conservation status:

Nationally Increasing (Robertson *et al.* 2021).

Regional conservation status: Regionally Critical (Crisp, 2020).

The kārearea / New Zealand falcon is the twelfth most frequently encountered native forest bird species in these five-minute bird counts. Kārearea / New Zealand falcons have only been detected on 19 occasions during these five-minute bird counts, providing too few detections to allow trends in encounter rates to be modelled. Kārearea / New Zealand falcon encounter rates have fluctuated from year to year between 2011 and 2022, from zero birds recorded in 2011, 2014 and 2019 to a mean of 0.03 birds encountered per bird count station in 2021. Despite these year-to-year

fluctuations, there does not appear to have been any overall trend in kārearea / New Zealand falcon encounter rates between 2011 and 2022 (Figure 3.26).

The kārearea / New Zealand falcon is the eleventh most frequently reported native forest bird species by citizen scientists in Wellington City, with 2,090 encounters recorded between 2011 and 2023 (Figure 3.27). These reports have a remarkably uniform distribution across the city, which suggests that these reports represent a relatively small number of highly mobile individuals ranging freely in native forest, suburban and urban habitats throughout the city. Several pairs of kārearea / New Zealand falcons are known to breed in Wellington City, at locations including Zealandia Te Māra a Tāne, Ōtari-Wilton's Bush, Tinakori Hill and on Miramar Peninsula (eBird, 2023). Kārearea / New Zealand falcon typically occur at fairly low densities and are known to be highly mobile (Heather & Robertson, 2015), so are not expected to undergo any dramatic increases in Wellington City in response to local conservation management efforts.




Figure 3.26: Mean (\pm SE) number of kārearea / New Zealand falcons recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.27: Distribution of karearea / New Zealand falcons in Wellington City between 2011 and 2023. Orange circles represent karearea / New Zealand falcon detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of karearea / New Zealand falcons detected per station between 2011 and 2022. Yellow and green circles represent karearea / New Zealand falcon observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.14 Toutouwai / North Island robin (*Petroica longipes*)

National conservation status: At Risk, Declining
(Robertson *et al.* 2021).

Regional conservation status: Not Threatened
(Crisp, 2020).

The toutouwai / North Island robin was reintroduced to Zealandia Te Māra a Tāne between 2001 and 2002 (Miskelly, 2018) and is now the fourteenth most frequently encountered native forest bird species in these five-minute bird counts. Toutouwai / North Island robins have only been detected on 14 occasions during these five-minute bird counts, providing too few detections to allow trends in encounter rates to be modelled. Toutouwai / North Island robin encounter rates have fluctuated from zero to a mean of 0.015 birds recorded per bird count each year, with no overall trend in encounter rates between 2011 and 2022 (Figure 3.28).



Image courtesy of Neil Fitzgerald/
NZ Birds Online

The toutouwai / North Island robin is the eighth most frequently reported native forest bird species by citizen scientists in Wellington City with 4,086 encounters recorded between 2011 and 2023, the majority of which are from within Zealandia Te Māra a Tāne. Of those reports from outside of Zealandia Te Māra a Tāne's predator-proof fence, the majority have occurred at locations within 500m of the fence, with only occasional records from forested reserves further afield, including in Ōtari-Wilton's Bush, Tinakori Hill, the Wellington Botanical Gardens, Prince of Wales Park and Makara Peak (Figure 3.29). This localised distribution of toutouwai / North Island robin observations in and around Zealandia Te Māra a Tāne suggests that one or more environmental factors are severely limiting the ability of this species to colonise forest habitats beyond Zealandia Te Māra a Tāne's predator-proof fence.




Figure 3.28: Mean (\pm SE) number of toutouwai / North Island robins recorded per five-minute bird count station in Wellington City between 2011 and 2022.




Figure 3.29: Distribution of toutouwai / North Island robins in Wellington City between 2011 and 2023. Orange circles represent toutouwai / North Island robin detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of toutouwai / North Island robins detected per station between 2011 and 2022. Yellow and green circles represent toutouwai / North Island robin observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.15 Tītipounamu / Rifleman (*Acanthisitta chloris*)



Image courtesy of Glenda Rees
NZ Birds Online

National Conservation status: At risk, Declining
(Robertson *et al.* 2021)

Regional conservation status: At risk, Declining
(Crisp, 2020).

Tītipounamu / rifleman apparently occurred in Wellington City until at least the mid-19th Century, as Oliver (1955) reports that the species was collected by P. Earle at Port Nicholson ‘about 1840’. Tītipounamu / rifleman were either extremely rare or extinct in the area by 1938 however, as Secker (1958) did not encounter them during extensive explorations made in the area between 1938 and 1956. Tītipounamu / rifleman were reintroduced to Zealandia Te Māra a Tāne in March 2019 when 59 birds were transferred from the Wainuiomata/Orongorongo Water Collection Area. These birds have bred successfully within Zealandia Te Māra a Tāne with 58 fledglings recorded during the 2020/2021 breeding season alone².

Tītipounamu / rifleman were detected for the first time during these five-minute bird counts in 2022, with a single bird being detected at a count station in Wright’s Hill Reserve 453 metres from the Zealandia Te Māra a Tāne fenceline. Tītipounamu / rifleman have also been reported on 281 occasions by citizen scientists since 2019, although the majority of these sightings have been made within Zealandia Te Māra a Tāne. A number of significant sightings have been made beyond Zealandia Te Māra a Tāne’s predator-proof fence however; for example during the 2021/2022 breeding season tītipounamu / rifleman bred successfully on Te Ahumairangi Hill and a family group was also encountered in Makara Peak Park, indicating that tītipounamu / rifleman have bred successfully at this latter location as well (Melissa Boardman *personal observation*; Thomas, 2022). The species has also been recorded recently in Central Park, Polhill Reserve and in Otari-Wilton Bush (eBird, 2023), providing an early indication that tītipounamu / rifleman are proving to be remarkably successful at colonising native forest habitats beyond Zealandia Te Māra a Tāne’s predator-proof fence (Figure 3.30).

² <https://www.facebook.com/ZEALANDIA/posts/10157605617536401>; accessed 04/04/2021.




Figure 3.30: Distribution of tītīpounamu / rifleman in Wellington City between 2011 and 2023. Orange circles represent tītīpounamu / rifleman detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of tītīpounamu / rifleman detected per station between 2011 and 2022. Yellow and green circles represent tītīpounamu / rifleman observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.16 Hihi (*Notiomystis cincta*)

National conservation status:

Nationally Vulnerable (Robertson *et al.* 2021).

Regional conservation status:

Regionally Critical (Crisp, 2020).

Hihi were reintroduced to Zealandia Te Māra a Tāne between 2005 and 2010, and a population is now being maintained in the sanctuary with an ongoing programme of supplementary feeding (Heather & Robertson, 2015; Miskelly, 2018). Hihi were detected during these five-minute bird counts for the first and only time in 2019, with a single bird being seen and heard at each of two count stations in Wrights Hill Reserve, 390 and 605 metres from the nearest point of Zealandia Te Māra a Tāne’s boundary fence respectively.



Image courtesy of Paul Le Roy/NZ Birds Online

Hihi is the fourteenth most frequently reported native forest bird species by citizen scientists in Wellington City, with 1,665 records submitted since 2011. The majority of these records have been from within Zealandia Te Māra a Tāne, however there have been several dozen reports of birds seen or heard in native forest habitats up to 600m from Zealandia Te Māra a Tāne’s boundary fence (Figure 3.31). In May 2016, a single hihi was reported from Orleans Recreational Reserve in Ngaio, 4.5 km NNE of Zealandia Te Māra a Tāne (Birds New Zealand, 2016). With this single exception, it appears that hihi are largely confined to Zealandia Te Māra a Tāne, and either experience very low emigration rates from the sanctuary, or very low survival rates once they leave the sanctuary.




Figure 3.31: Distribution of hihi in Wellington City between 2011 and 2023. Orange circles represent hihi detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of hihi detected per station between 2011 and 2022. Yellow circles represent hihi observations reported by local citizen scientists via eBird between 2011 and 2023.

3.2.17 Miromiro / Tomtit (*Petroica macrocephala*)

National conservation status: Not Threatened
(Robertson *et al.* 2021).

Regional conservation status: Not Threatened
(Crisp, 2020).

The miromiro / tomtit is a vagrant (i.e., irregular visitor) to Wellington City at the present time, with no local self-sustaining population known to currently exist within Wellington City boundaries. A single miromiro / tomtit was recorded for the first time during this five-minute bird count project in 2016, at a count station in Khandallah Park (Figure 3.32; McArthur *et al.* 2017). In January 2021, at least one tomtit was recorded in Ōtari-Wilton's Bush by New Zealand Bird Atlas surveyors (Milius, 2021). The nearest miromiro / tomtit breeding populations to Wellington City currently occur in Belmont and East Harbour Regional Parks, approximately 13 and 14 km from Wellington City respectively (eBird, 2023). Miromiro / tomtits are known to have strong dispersal abilities, including over open water (Parker *et al.* 2004), so it is likely that these recent Wellington City observations represent birds that have dispersed into the city from one of these nearby source populations. Several more recent citizen science observations recorded in Zealandia may be misidentifications of similar species (Figure 3.32).

Miromiro / tomtits were present in native forest habitats within Wellington City limits historically and persisted in both Ōtari-Wilton's Bush and Khandallah Park until at least as late as the mid-1920s (Stidolph, 1924; 1925). An attempt was made to reintroduce miromiro / tomtits to Zealandia Te Māra a Tāne between 2001 and 2004 and some successful local breeding subsequently occurred both inside and outside of Zealandia's predator proof-fence between 2003 and 2006. However, this population suffered from high rates of emigration, possibly caused by interspecific competition with an already well-established toutouwai / North Island robin population. As a result, miromiro / tomtits failed to establish in Zealandia Te Māra a Tāne and had become locally extinct again by 2007 (Empson and Fastier, 2013).



Image courtesy of Paul Shaw/NZ Birds Online




Figure 3.32: Distribution of miromiro / tomtits in Wellington City between 2011 and 2023. Orange circles represent miromiro / tomtit detections at five-minute bird count stations, with the size of the circle corresponding to the mean annual number of miromiro / tomtits detected per station between 2011 and 2022. Yellow and green circles represent miromiro / tomtit observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.2.18 Koekoeā / Long-tailed cuckoo (*Eudynamys taitensis*)



Image courtesy of Adam Clarke/NZ Birds Online

National conservation status: Nationally Vulnerable (Robertson *et al.* 2021).

Regional conservation status: At Risk, Naturally Uncommon (Crisp, 2020).

Koekoeā / long-tailed cuckoos appear to be a rare passage migrant in Wellington City at the present time. Koekoeā / long-tailed cuckoos have not yet been recorded during these five-minute bird counts and have only been recorded by citizen scientists in Wellington City on twelve occasions since 2011 (Figure 3.33). Given the current scarcity of koekoeā / long-tailed cuckoos in Wellington City, the local population of

pōpokotea / whiteheads present in forest habitats in and adjacent to Zealandia Te Māra a Tāne is likely to be the only pōpokotea / whitehead population in the Wellington region that does not experience brood parasitism by koekoeā / long-tailed cuckoos.

3.2.19 Ruru / Morepork (*Ninox novaeseelandiae*)



Image courtesy of Adam Clarke/
NZ Birds Online

National conservation status: Not Threatened (Robertson *et al.* 2021).

Regional conservation status: Not Threatened (Crisp, 2020).

The ruru / morepork is the most common and conspicuous native forest bird species present in Wellington City that has not yet been detected during these five-minute bird counts, owing to its nocturnal habits and the fact that these counts are carried out during daylight hours. They have been reported on 979 occasions in Wellington City by citizen scientists since 2011. These records suggest that ruru / moreporks are relatively widespread in Wellington City but are largely restricted to the larger forested reserves forming Wellington's 'green belt' including Zealandia Te Māra a Tāne, and to the eastern parts of Miramar Peninsula (Figure 3.34).




Figure 3.33: Distribution of koekoeā / long-tailed cuckoos in Wellington City between 2011 and 2023. Orange circles represent koekoeā / long-tailed cuckoo non-detections at five-minute bird count stations between 2011 and 2022. Yellow circles represent koekoeā / long-tailed cuckoo observations reported by local citizen scientists via eBird between 2011 and 2023.




Figure 3.34: Distribution of ruru / moreporks in Wellington City between 2011 and 2023. Orange circles represent ruru / morepork non-detections at five-minute bird count stations between 2011 and 2022. Yellow and green circles represent ruru / morepork observations reported by local citizen scientists via eBird and iNaturalist between 2011 and 2023.

3.3 The Zealandia Te Māra a Tāne ‘halo effect’

The best model to describe how the proximity of a bird count station to Zealandia Te Māra a Tāne influences the diversity of native and introduced species at that station was the model that assumed different spatial trends for native versus introduced species. This model had a lower AICc value ($\Delta\text{AICc} = 6.54$) than the next best model, which was the model that assumed that native and introduced species would show the same spatial trend.

According to this model, the species richness of native birds was higher at count stations closer to Zealandia Te Māra a Tāne, suggesting that the predator-free sanctuary is exerting a measurable ‘halo’ effect on forest bird communities beyond the predator-proof fence (the coefficient of distance on log number of native forest bird species was -0.577 ; 95% CI $-0.620 - -0.534$). On average, a mean of $6.8 (\pm 0.3 \text{ SE})$ native forest bird species were detected at count stations closest to Zealandia Te Māra a Tāne’s boundary, whereas a mean of $4.6 (\pm 0.3 \text{ SE})$ native forest bird species were detected at the stations situated farthest away from Zealandia Te Māra a Tāne (a distance of approximately 10 km).

In contrast, the species richness of introduced bird species increased slightly with increasing distance from Zealandia Te Māra a Tāne (the coefficient of distance on log number of introduced bird species was 0.011 ; 95% CI $-0.022 - 0.038$). On average, a mean of $2.1 (\pm 0.3 \text{ SE})$ introduced bird species were detected at count stations closest to Zealandia Te Māra a Tāne’s boundary, whereas a mean of $3.5 (\pm 0.5 \text{ SE})$ introduced species were detected at the stations situated farthest away from Zealandia Te Māra a Tāne (Figure 3.35).




Figure 3.35: Mean number of native and introduced bird species detected per station between 2011 and 2023, in relation to each station’s distance from Zealandia Te Māra a Tāne.

4. Discussion

4.1 Changes in bird community composition, abundance and distribution

The mean number of native forest birds being counted per five-minute bird count has increased by 41% since 2011, and this has been driven by increases in the mean number of tūī, pīwakawaka / New Zealand fantail, kākā, kererū and tīeke / North Island saddleback being counted each year. Two of these species were absent from Wellington City prior to being reintroduced to the predator-free Zealandia Te Māra a Tāne sanctuary during the early 2000s, and all of these species are vulnerable to depredation by mammalian predators to varying degrees (Innes *et al.* 2010; Fea & Hartley, 2018). Given that these five-minute bird counts are carried out at the same time each year, in the same weather conditions and usually by the same observers, these results provide strong evidence that the abundance of these five species has increased in Wellington City since 2011. This being the case, it is clear that these increases in native forest bird abundance are being driven by both the establishment of large source populations of these species within Zealandia Te Māra a Tāne, and the ongoing expansion and intensification of mammalian predator control efforts across Wellington City.

Although mean annual counts for native forest birds have increased in native forest habitats in Wellington City parks and reserves between 2011 and 2022, mean annual counts for introduced species have also increased at a similar rate. This suggests that the native forest bird communities in forested parks and reserves outside of Zealandia Te Māra a Tāne have not yet recovered to the point where interspecific competitive exclusion is reducing the abundance and conspicuousness of introduced bird species, as has now occurred within Zealandia Te Māra a Tāne itself (Miskelly, 2018). One local exception to this pattern currently exists, however. Mean annual counts of introduced birds at five-minute bird count stations within a few hundred metres of Zealandia Te Māra a Tāne's predator-proof fence are lower than those stations situated furthest away from this predator-free sanctuary. This indicates that the interspecific competitive exclusion of introduced bird species that is occurring within Zealandia Te Māra a Tāne is also extending some distance beyond Zealandia Te Māra a Tāne's predator-proof fence, into surrounding forest habitats in adjacent parks and reserves, part of what is colloquially known as Zealandia Te Māra a Tāne's 'halo' effect.

Zealandia Te Māra a Tāne's 'halo' effect is not confined to introduced species, however. On average, almost twice as many native bird species have been recorded annually at those five-minute bird count stations situated closest to Zealandia Te Māra a Tāne's boundary than at those situated furthest away. These higher levels of native bird species richness at stations in proximity to Zealandia Te Māra a Tāne are being driven by the emigration of birds from the large source populations now established within Zealandia Te Māra a Tāne. This measurable 'halo' effect demonstrates the key role that Zealandia Te Māra a Tāne is playing in driving increases in the diversity, abundance and distribution of native forest birds in Wellington City's parks and reserves.

This in turn raises the question of the relative contributions that Zealandia Te Māra a Tāne's 'halo' effect and city-wide mammalian predator control initiatives are making towards the improvements in native forest bird populations observed in Wellington City since 2011. Brudvig *et al.* (2009) for instance report that biodiversity 'spillover' in a landscape is largely a function of processes occurring within the source population rather than in the surrounding landscape, suggesting that some of the increases in native forest bird diversity, abundance and distribution observed in Wellington City may have

occurred irrespective of the extent or intensity of mammalian predator control occurring beyond Zealandia Te Māra a Tāne's predator-proof fence. Indeed, it has long been recognised that species occupancy and density is a poor predictor of habitat quality, and that even relatively poor-quality habitats can support high population densities due to immigration alone (e.g., McArthur *et al.* 2019).

In the case of Wellington City, the distribution maps contained in this report suggest that the relative importance of conservation management initiatives both within and beyond Zealandia Te Māra a Tāne's predator-proof fence is likely to be species specific. Based on these distribution maps, Wellington City's resident native forest bird species can be divided into two broad categories: those species with relatively widespread distributions, and those with very localised distributions. Several widespread species, including tūi, pīwakawaka / New Zealand fantail and kererū, are now frequently observed breeding successfully in native forest and suburban habitats beyond Zealandia Te Māra a Tāne's predator-proof fence, so are more likely to be benefitting from city-wide reductions in mammalian predator densities in addition to being supplemented by immigration from Zealandia Te Māra a Tāne.

A number of other species continue to have very localised distributions centred on Zealandia Te Māra a Tāne, including toutouwai / North Island robin, and tīeke / North Island saddleback. The presence of these species in native forest habitats adjacent to Zealandia Te Māra a Tāne appears to be largely a result of the emigration of birds from Zealandia Te Māra a Tāne, rather than of improvements in local predator control efforts in the reserves in which these birds are being observed. For example, toutouwai / North Island robins have been well established in Zealandia Te Māra a Tāne for at least 15 years (McGavin, 2009; Empson & Fastier, 2013), yet have only been detected on 14 occasions during these five-minute bird counts between 2011 and 2022 and are seldom reported by citizen scientists at distances greater than around 1 km from Zealandia Te Māra a Tāne (Figure 3.29). New Zealand robins (*Petroica* spp.) are known to have relatively strong dispersal capabilities through habitats dominated by woody vegetation, with juvenile birds capable of dispersing up to 11 km from their natal territories in forested habitat (Oppel & Beaven, 2004; Richard, 2007), so habitat connectivity is unlikely to be the factor limiting the expansion of this species in Wellington City. Mark-resighting and nest monitoring of toutouwai / North Island robins in reserves adjacent to Zealandia in recent years has confirmed that although these reserves provide suitable breeding habitat for robins, poor adult and juvenile survival rates are limiting the ability of this species to colonise forest habitats outside of Zealandia Te Māra a Tāne. Of 37 adult toutouwai / North Island robins banded in forested reserves adjacent to Zealandia Te Māra a Tāne in the winter of 2017, only 11 birds were still present on their territories by the beginning of the following breeding season, and only four were still present in the winter of 2018. During the 2017-18 breeding season, a total of ten nesting attempts were monitored. Eight of these successfully fledged young, with 11 young subsequently reaching independence. However, only three of these 11 offspring were detected subsequently, and only one appeared to survive long enough to attempt to breed (unsuccessfully) the following season. Of the two nests that failed, one was depredated by a cat, and the other by a stoat (Shaw and Harvey, 2018). Adult survival rates were similarly low during the 2018-19 breeding season. Of 16 adult birds known to be present on breeding territories in September 2018, only one was still present by March 2019. Breeding success during this second season was substantially worse than the previous year. Of seven nests monitored during the 2018-19 season, only one successfully fledged young (MacKinlay, unpublished data).

Given this evidence, the most likely factor limiting the establishment of toutouwai / North Island robins (as well as other species with localised Wellington City distributions, including tīeke / North Island saddleback and kākāriki / red-crowned parakeet) beyond the boundaries of Zealandia Te Māra

a Tāne is depredation of both adult and juvenile birds by mammalian predators, particularly both domestic and wild cats (*Felis catus*) and mustelids. Evidence is growing that cats are now the most significant predator of native birds in Wellington City. For example, of 23 kākārīki / red-crowned parakeets that were recovered dead outside of Zealandia Te Māra a Tāne's predator-proof fence between 2010 and 2016, nine of these birds (39%) had been depredated by cats (Irwin & Empson, 2022). Although considerable effort is being invested in reducing populations of rats, possums and mustelids, cats are currently not being targeted. Camera trapping work carried out by researchers at Victoria University of Wellington has shown that cats accounted for a relatively large proportion of the approximately 22,000 animal 'detections' collected from several Wellington City reserves over a five-month period in 2014 (<http://identifyanimals.co.nz/>; accessed 24/09/2015; Anton *et al.* 2018), suggesting that cats occur in relatively high densities in the parks and reserves that were sampled. Further camera trapping work carried out in 2016 confirmed that cats were likely to be present across most of the total area of Polhill Reserve (one of the forested reserves adjoining predator-free Zealandia Te Māra a Tāne), and that the majority of these cats appeared to be domestic pets (Woolley & Hartley, 2019).

Given this growing body of evidence, we agree with the conclusion drawn by Shaw & Harvey (2018), that if Wellington City Council and Predator Free Wellington wish to create "a natural city that flourishes with native wildlife and a dawn chorus that will be the envy of other cities" (<https://www.pfw.org.nz/>; accessed 25/06/2019), then significant work will need to be done to manage the risk to wildlife posed by feral, stray and free-roaming domestic cats in Wellington City. Until this occurs, further increases in the diversity, abundance and distribution of native forest birds in Wellington City will be constrained by the city's large cat population. Specifically, creating healthy, productive populations of native forest birds such as toutouwai / North Island robin, tieke / North Island saddleback, pōpokotea / whitehead and kākārīki / red-crowned parakeets in forest habitats outside of Zealandia Te Māra a Tāne's predator-proof fence is unlikely to be attainable given the presence of a large cat population, irrespective of the degree to which other mammalian predators such as rats, possums, mustelids and hedgehogs are controlled or eradicated.

4.2 The role of citizen scientists in monitoring Wellington City's bird fauna

Citizen scientists are playing an increasingly important role in providing bird observation data that complement this Wellington City five-minute bird count dataset, enabling us to map the distribution of birds in Wellington City to a level of detail never done before. A total of 95,761 verified observations of native forest birds have been contributed by citizen scientists in Wellington City between 2011 and 2023 and are included on the distribution maps in this report. Ninety-three percent or 89,104 of these observations have been contributed via the New Zealand eBird database, making eBird by far the most popular database used by Wellington-based citizen scientists that have an interest in birds. A further 6% (6,204 observations) were submitted via the iNaturalist New Zealand database, making this the second most preferred database used by Wellington-based citizen scientists. An additional 1% of records (574 observations) were sourced from the Great Kererū Count.

The contribution that citizen scientists have made to this bird monitoring programme has provided at least two benefits. Firstly, because citizen scientists are collecting observations from a large number of locations not being sampled as part of this project, the resulting bird distribution maps are considerably more detailed than if they had been derived from the five-minute bird count data alone. These distribution maps have provided insights into the habitat use of native bird species in Wellington

City and have greatly improved our ability to map the steadily expanding ranges of species such as kākā and kākārīki / red-crowned parakeets as they colonise native forest habitats beyond Zealandia Te Māra a Tāne's predator-proof fence. The second benefit is the role that citizen scientists play in detecting rare, cryptic or re-colonising bird species in Wellington City. Citizen scientists are much better placed to detect colonising species or those species which are naturally rare, cryptic or nocturnal because they are collecting bird observations year-round. Indeed, our knowledge of the presence and distribution of two native forest bird species that occur in Wellington City (ruru / morepork and koekoeā / long-tailed cuckoo), is entirely derived from citizen science data at present, as none of these species have yet been detected during a five-minute bird count.

Although our knowledge of the abundance and distribution of diurnal, or day-active, bird species in Wellington City has improved substantially over the past twelve years that these five-minute bird counts have been underway, the abundance and distribution of the one relatively widespread nocturnal forest bird species present in Wellington City is very poorly understood. Ruru / morepork may well be relatively common in Wellington City, and trends in ruru / morepork encounter rates or distribution over time could provide an additional measure of the outcomes of local pest control efforts. An opportunity exists therefore to fill this knowledge gap by running a citizen science project specifically aimed at mapping the distribution of ruru / morepork in Wellington City and quantifying encounter rates as an indirect measure of abundance. Such a project could be modelled on the 2011 Hamilton City ruru / morepork survey, whereby volunteers were assigned to a pre-defined set of survey locations over a period of five consecutive nights (Morgan & Styche, 2012). This project would also serve a secondary purpose of providing Wellington City residents with an additional opportunity to engage with their surrounding natural environment, learn more about the birds around them and improve their skills as citizen scientists.

5. Recommendations

Based on the results described in this report, we suggest that Wellington City Council considers adopting the following recommendations:

- That this five-minute bird count monitoring programme be continued on an annual basis, to provide a consistent, repeatable measure of the state and trends in the diversity, distribution and abundance of birds in Wellington City parks and reserves, in order to contribute towards objective 4.2.2a of WCC's Biodiversity Strategy and Action Plan (WCC, 2015).
- That local citizen scientists continue to be encouraged to contribute their bird observations by submitting them in the form of complete bird checklists to the New Zealand eBird database (objective 4.3.3a of WCC's Biodiversity Strategy and Action Plan; WCC, 2015). By doing so, local citizen scientists will be submitting their Wellington City bird observation data to a single, centralised database, providing researchers and conservation managers with easy access to a large and fast-growing dataset of high quality Wellington City bird observations.
- That Wellington City residents continue to be educated about the risk that domestic cats pose to the city's wildlife, and options continue to be explored to manage the risk that domestic, stray and feral cats pose to native and endemic birds in Wellington City.
- That a citizen science project be carried out with the aim of mapping the distribution of ruru / morepork in Wellington City during the summer of 2023/2024 (objective 3.3.4b of WCC's Biodiversity Strategy and Action Plan; WCC, 2015). Such a project could involve public requests for ruru / morepork sightings during a pre-defined time period (e.g., the month of November), coupled with recruiting a pool of local volunteers to carry out night-time surveys of a pre-determined network of locations throughout the city to determine ruru / morepork distribution in local parks and reserves.

Acknowledgements

This work was instigated by Myfanwy Emeny, Open Space and Parks Manager at Wellington City Council and by Philippa Crisp, former Team Leader - Land, Ecology and Climate at Greater Wellington Regional Council. We would like to thank Roger Uys at Greater Wellington Regional Council for the administrative and logistical support he has provided to this project, and Mark McAlpine for assisting in carrying out the 2011 counts.

We also extend our thanks to Raewyn Empson (former Conservation Manager at Zealandia Te Māra a Tāne) for providing us with access to bird distribution data held by the Karori Sanctuary Trust, and to Tony Stoddard (Kererū Discovery) for providing us with a copy of the Great Kererū Count citizen science dataset. Our thanks also to Tim Park (Wellington City Council) for his assistance with downloading citizen science data from iNaturalist New Zealand. We would also like to acknowledge the valuable efforts of a growing number of dedicated and observant citizen scientists who are collecting and sharing their Wellington City bird observations via the New Zealand eBird and iNaturalist New Zealand databases.

Thanks also to Luke Crouch, Richard Farrell, Li Loo, Samantha Ray, Grant Redvers, and Roger Uys for assisting with data entry and management, and to Anne Schlesselmann and Adrian Monks at Manaaki Whenua Landcare Research for assisting with the statistical analyses included in this report. We are also grateful to Rebecca Lunt and Roger Uys for reviewing earlier drafts of this report.

References

- Anton, V.; Hartley, S. and Wittmer, H. 2018. Evaluation of remote cameras for monitoring multiple invasive mammals in New Zealand. *New Zealand Journal of Ecology* 42: 74-79.
- Bates, D.; Mächler, M.; Bolker, B. and Walker, S. 2015. Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software* 67: 1–48.
- Bell, B.D. 2008. Tūi (*Prosthemadera novaeseelandiae*) increase at Seatoun, Miramar Peninsula, Wellington, New Zealand during 1998-2006. *Notornis* 55: 104-106.
- Bibby, C.J.; Burgess, N.D.; Hill, D.A. and Mustoe, S. 2000. *Bird census techniques* (2nd edition). Academic Press, London.
- Birds New Zealand, 2016. eBird checklist: <https://ebird.org/newzealand/checklist/S30471770>. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: 4th April, 2021).
- Brockie, R.E. and Duncan, C. 2012. Long term trends in Wellington City bird counts: 1969-2006. *Notornis* 59: 1-6.
- Brudvig, L.A.; Damschen, E.I.; Tewksbury, J.J.; Haddad, N.M.; Levey, D.J. 2009. Landscape connectivity promotes plant biodiversity spillover into non-target habitats. *Proceedings of the National Academy of Sciences of the United States of America* 106: 9328–9332. doi: 10.1073/pnas.0809658106.
- Burnham, K.P. and Anderson, D.R. 2002. *Model selection and multimodel inference: A practical information-theoretic approach*. Springer, New York.
- Checklist Committee (OSNZ). 2022. *Checklist of the Birds of New Zealand* (5th edition). Ornithological Society of New Zealand Occasional Publication No. 1. Ornithological Society of New Zealand, Wellington.
- Crisp, P. 2020. *Conservation status of native bird species in the Wellington region*. Publication No. GW/ESCI-G-20/75, Greater Wellington Regional Council, Wellington.
- Dawson, D.G. and Bull, P.C. 1975. Counting birds in New Zealand forests. *Notornis* 22: 101-109.
- eBird, 2023. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: 20/04/2023).
- Empson, R. and Fastier, D. 2013. Translocations of North Island tomtits (*Petroica macrocephala toitoi*) and North Island robins (*P. longipes*) to Zealandia-Karori Sanctuary, an urban sanctuary. What have we learned? *Notornis* 60: 63-69.
- Fea, N. and Hartley, S. 2018. The balancing act of nest survival: survival of a small endemic bird in the face of ship rat predation and other risk factors. *Avian Conservation and Ecology* 13(2):11. <https://doi.org/10.5751/ACE-01284-130211>.
- Froude, V.A. 2009. *Changes in native forest bird distribution and abundance in Wellington City Council reserves 2001-2009*. Unpublished report for Wellington City Council, Pacific Eco-Logic Ltd, Russell.

- Gelman, A. and Hill, J. 2007. *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press, New York.
- Harrison, X.A. 2014. Using observation-level random effects to model overdispersion in count data in ecology and evolution. *PeerJ* 2:e616 <https://doi.org/10.7717/peerj.616>.
- Hartley, L. and Greene, T. 2012. Incomplete counts: Five-minute bird counts. DoC inventory and monitoring toolbox (DOCDM-534972), Department of Conservation, Wellington. (<http://www.doc.govt.nz/Documents/science-and-technical/inventory-monitoring/im-toolbox-birds-incomplete-five-min-counts.pdf>; accessed 05/04/2021).
- Hartley, S. 2021. *Great Kererū Count 2021 Annual report*. Unpublished report, Centre for Biodiversity and Restoration Ecology, Victoria University, Wellington.
- Heather, B.D. and Robertson, H.A. 2015. *The field guide to the birds of New Zealand*, 4th edition. Penguin, Auckland.
- Innes, J.; Kelly, D.; McC. Overton, J. and Gillies, C. 2010. Predation and other factors currently limiting New Zealand forest birds. *New Zealand Journal of Ecology* 34: 86-114.
- Irwin, E.T. and Empson, R.A. 2022. Red-crowned parakeet (kākāriki, *Cyanoramphus novaezelandiae*) mortality inside and outside a fenced sanctuary in Wellington City, New Zealand. *Notornis* 69: 54-58.
- Mackenzie, D.I.; Nichols, J.D.; Royle, J.A.; Pollock, K.H.; Bailey, L.L. and Hines, J.E. 2006. *Occupancy estimation and modelling: inferring patterns and dynamics of species occurrence*. Elsevier Academic Press, Burlington, USA.
- McArthur, N.; Boulton, R.L.; Richard, Y.; Armstrong, D.P. 2019. The role of pine plantations in source-sink dynamics of North Island robins. *New Zealand Journal of Ecology* 43: 3362. doi: 10.20417/nzj ecol.43.12
- McArthur, N.; Flux, I. and Harvey, A. 2017. *State and trends in the diversity, abundance and distribution of birds in Wellington City*. Client report prepared for Greater Wellington Regional Council. Wildlife Management International Ltd, Blenheim.
- McArthur, N.; Flux, I. and Harvey, A. 2021. *State and trends in the diversity, abundance and distribution of birds in Wellington City*. Client report prepared for Greater Wellington Regional Council, Wellington.
- McArthur, N.; Flux, I. and Harvey, A. 2022. *State and trends in the diversity, abundance and distribution of birds in Wellington City*. Client report prepared for Greater Wellington Regional Council, Wellington.
- McArthur, N.; Harvey, A. and Flux, I. 2013. *State and trends in the diversity, abundance and distribution of birds in Wellington City reserves*. Greater Wellington Regional Council, Publication No. GW/ESCI-T-14/43, Wellington.
- McArthur, N.; Moylan, S. and Crisp, P. 2012. *Baseline survey of the diversity, abundance and distribution of birds in Wellington City reserves, June 2012*. Greater Wellington Regional Council, Publication No. GW/EMI-T-12/231, Upper Hutt.
- McGavin, S. 2009. Density and pair fidelity in a translocated population of North Island robin (*Petroica longipes*). *Notornis* 56: 206-212.

- McLaughlin, M. and Harvey, A. 2013. *Bellbird (Anthornis melanura) korimako translocation report, March 2013*. Unpublished report, Friends of Mana Island, Wellington.
- Milius, N. 2021. eBird checklist: <https://ebird.org/atlasnz/checklist/S78842614>. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: 4th April, 2021).
- Miskelly, C.M.; Empson, R. and Wright, K. 2005. Forest birds recolonising Wellington. *Notornis* 52: 21-26.
- Miskelly, C.M. and Powlesland, R.G. 2013. Conservation translocations of New Zealand birds 1863-2012. *Notornis* 60: 3-28.
- Miskelly, C.M. 2018. Changes in the forest bird community of an urban sanctuary in response to pest mammal eradications and endemic bird reintroductions. *Notornis* 65: 132-151.
- Morgan, D.K.J. and Styche, A. 2012. Results of a community-based acoustic survey of ruru (moreporks) in Hamilton city. *Notornis* 59: 123-129.
- Oppel, S. and Beaven, B. 2004. Juvenile Stewart Island robins (*Petroica australis rakiura*) disperse up to 16 km. *Notornis* 51: 55-56.
- Oliver, W.R.B. 1955. *New Zealand Birds*. A.H. and A.W. Reed, Wellington.
- Parker, K.A.; Hughes, B.; Thorogood, R. and Griffiths, R. 2004. Homing over 56 km by a North Island tomtit (*Petroica macrocephala toitoi*). *Notornis* 51: 238-239.
- R Core Team 2021. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org>.
- Richard, Y. 2007. *Demography and distribution of the North Island robin (Petroica longipes) in a fragmented agricultural landscape of New Zealand*. Doctoral dissertation, Massey University, Palmerston North, New Zealand.
- Robertson, H.A.; Baird, K.A.; Elliott, G.P.; Hitchmough, R.A.; Miskelly, C.M.; McArthur, N.J.; Makan, T.D.; Miskelly, C.M.; O'Donnell, C.J.F.; Sagar, P.M.; Scofield, R.P.; Taylor, G.A. and Michel, P. 2021. *Conservation status of New Zealand birds, 2021*. New Zealand Threat Classification Series 36, Department of Conservation, Wellington.
- Scofield, R.P.; Christie, D.; Sagar, P.M. and Sullivan, B.L. 2012. eBird and avifaunal monitoring by the Ornithological Society of New Zealand. *New Zealand Journal of Ecology* 36: 279-286.
- Secker, H.L. 1958. Status of Rifleman in hillside scrub land of Wellington district. *Notornis* 8: 23.
- Shaw, R. and Harvey, A. 2018. *Toutouwai population monitoring and community engagement in WCC reserves adjoining Zealandia, 2017-2018*. Unpublished report submitted to Wellington City Council, Victoria University Wellington, Wellington.
- Stidolph, R.H.D. 1924. eBird checklist: <http://ebird.org/ebird/newzealand/view/checklist/S21360197>. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: 30th June 2017).
- Stidolph, R.H.D. 1925. eBird checklist: <http://ebird.org/ebird/newzealand/view/checklist/S31000943>. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: 30th June 2017).

Sullivan, B.L.; Aycrigg, J.L.; Barry, J.H.; Bonney, R.E.; Bruns, N.; Cooper, C.B.; Damoulas, T.; Dhondt, A.A.; Dietterich, T.; Farnsworth, A.; Fink, D.; Fitzpatrick, J.W.; Fredericks, T.; Gerbracht, J.; Gomes, C.; Hochachka, W.M.; Iliff, M.J.; Lagoze, C.; La Sorte, F.; Merrifield M.; Morris, W.; Phillips, T.B.; Reynolds, M.; Rodewald, A.D.; Rosenberg, K.V.; Trautmann, N.M.; Wiggins, A.; Winkler, D.W.; Wong, W.-K.; Wood, C.L.; Yu, J. and Kelling, S. 2014. The eBird enterprise: An integrated approach to development and application of citizen science. *Biological Conservation* 169: 31-40.

Thomas, D. 2022. eBird checklist: <https://ebird.org/atlasnz/checklist/S100446709>. eBird: An online database of bird distribution and abundance [web application]. eBird, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: 28th April, 2022).

Wellington City Council, 2015. *Our natural capital – Wellington’s biodiversity strategy and action plan 2015*. Wellington City Council, Wellington.

Woolley, C.K. and Hartley, S. 2019. Activity of free-roaming domestic cats in an urban reserve and public perception of pet-related threats to wildlife in New Zealand. *Urban Ecosystems* <https://doi.org/10.1007/s11252-019-00886-2>.

Appendix One

This appendix contains a list of all bird species encountered in Wellington City parks and reserves during five-minute bird counts carried out between 2011 and 2022 (P = species detected). Scientific names, common names (both Māori and English) and the taxonomic order in which the species appear in this list all follow those used in the fifth edition of the *Checklist of the Birds of New Zealand* (Checklist Committee (OSNZ), 2022). The national conservation status rankings used are those New Zealand Threat Classification System rankings listed in Robertson *et al.* (2021) and the regional conservation status rankings are those listed in Crisp (2020).

Māori Name	Common Name	Scientific Name	National Conservation Status	Regional Conservation Status	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
pūtangitangi	paradise shelduck	<i>Tadorna variegata</i>	Not Threatened	Not Threatened		P	P				P	P				
rakiraki	mallard	<i>Anas platyrhynchos</i>	Introduced and Naturalised	Introduced and Naturalised					P	P						
koera	California quail	<i>Callipepla californica</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
heihei	feral chicken	<i>Gallus gallus</i>	Not Applicable ³	Not Applicable ³	P	P	P	P	P	P	P			P	P	

³ Although feral chickens are known to breed in the wild in New Zealand, no viable populations have been established and existing populations are being maintained by ongoing releases and/or supplementary feeding (Heather & Robertson, 2015). Because of this, the feral chicken is not recognised as a naturalised species in New Zealand (Checklist Committee (OSNZ) 2022) and therefore does not have a New Zealand Threat Classification System ranking (Robertson *et al.* 2021).

Māori Name	Common Name	Scientific Name	National Conservation Status	Regional Conservation Status	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
peihana	common pheasant	<i>Phasianus colchicus</i>	Introduced and Naturalised	Introduced and Naturalised											P	
kererū aropari	rock pigeon	<i>Columba livia</i>	Introduced and Naturalised	Introduced and Naturalised						P	P					P
kererū	kererū	<i>Hemiphaga novaeseelandiae</i>	Not Threatened	At Risk, Recovering	P	P	P	P	P	P	P	P	P	P	P	P
pīpīwharau	shining cuckoo	<i>Chrysococcyx lucidus</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P
tōrea pango	variable oystercatcher	<i>Haematopus unicolor</i>	At Risk, Recovering	Regionally Vulnerable		P		P	P	P		P	P		P	P
	spur-winged plover	<i>Vanellus miles</i>	Not Threatened	Not Threatened				P				P				
tarāpunga	red-billed gull	<i>Larus novaehollandiae</i>	At Risk, Declining	Regionally Vulnerable		P	P					P		P	P	P
karoro	southern black-backed gull	<i>Larus dominicanus</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P
tara	white-fronted tern	<i>Sterna striata</i>	At Risk, Declining	Regionally Endangered								P		P		P
tākupu	Australasian gannet	<i>Morus serrator</i>	Not Threatened	Migrant												P
kororā	little penguin	<i>Eudyptula minor</i>	At Risk, Declining	Regionally Vulnerable							P					
kawaupaka	little shag	<i>Phalacrocorax melanoleucos</i>	At Risk, Relict	Regionally Vulnerable										P		

Māori Name	Common Name	Scientific Name	National Conservation Status	Regional Conservation Status	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
kawau	black shag	<i>Phalacrocorax carbo</i>	At Risk, Relict	Regionally Critical								P				
kāruhiruhi	pied shag	<i>Phalacrocorax varius</i>	At Risk, Recovering	Regionally Vulnerable					P							
kawau tikitiki	spotted shag	<i>Stictocarbo punctatus</i>	Nationally Vulnerable	Regionally Endangered							P					
matuku moana	white-faced heron	<i>Egretta novaehollandiae</i>	Not Threatened	Not Threatened			P									
kāhu	swamp harrier	<i>Circus approximans</i>	Not Threatened	Not Threatened	P	P		P								P
kōtare	New Zealand kingfisher	<i>Todiramphus sanctus</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P
kārearea	New Zealand falcon	<i>Falco novaeseelandiae</i>	Nationally Increasing	Regionally Critical		P	P		P	P	P	P		P	P	P
kākā	kākā	<i>Nestor meridionalis</i>	At Risk, Recovering	At Risk, Recovering	P	P	P	P	P	P	P	P	P	P	P	P
kākā uhi whero	eastern rosella	<i>Platycercus eximius</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
kākāriki	red-crowned parakeet	<i>Cyanoramphus novaeseelandiae</i>	At Risk, Relict	At Risk, Recovering	P	P		P	P	P	P	P	P	P	P	P
tītītipounamu	rifleman	<i>Acanthisitta chloris</i>	At Risk, Declining	At Risk, Declining												P
korimako	bellbird	<i>Anthornis melanura</i>	Not Threatened	Not Threatened		P	P	P		P	P	P	P	P	P	P
tūī	tūī	<i>Prothemadera novaeseelandiae</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P

Māori Name	Common Name	Scientific Name	National Conservation Status	Regional Conservation Status	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
riroriro	grey warbler	<i>Gerygone igata</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P
tīeke	North Island saddleback	<i>Philesturnus rufusater</i>	At Risk, Recovering	Regionally Vulnerable	P	P	P	P	P	P	P	P	P	P	P	P
hihi	stitchbird	<i>Notiomystis cincta</i>	Nationally Vulnerable	Regionally Critical									P			
pōpokotea	whitehead	<i>Mohoua albicilla</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P
makipai	Australian magpie	<i>Gymnorhina tibicen</i>	Introduced and Naturalised	Introduced and Naturalised	P		P	P	P	P	P	P		P	P	P
pīwakawaka	New Zealand fantail	<i>Rhipidura fuliginosa</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P
miromiro	tomtit	<i>Petroica macrocephala</i>	Not Threatened	Not Threatened						P						
toutouwai	North Island robin	<i>Petroica longipes</i>	At Risk, Declining	Not Threatened	P	P	P		P			P	P		P	P
kairaka	skylark	<i>Alauda arvensis</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P		P	P	P	P	P	P
warou	welcome swallow	<i>Hirundo neoxena</i>	Not Threatened	Not Threatened	P					P	P					P
tauhou	silvereve	<i>Zosterops lateralis</i>	Not Threatened	Not Threatened	P	P	P	P	P	P	P	P	P	P	P	P

Māori Name	Common Name	Scientific Name	National Conservation Status	Regional Conservation Status	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
tāringi	common starling	<i>Sturnus vulgaris</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
manu pango	Eurasian blackbird	<i>Turdus merula</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
manu-kai-hua-rakau	song thrush	<i>Turdus philomelos</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
	dunnock	<i>Prunella modularis</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
tiu	house sparrow	<i>Passer domesticus</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
pahirini	chaffinch	<i>Fringilla coelebs</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
	greenfinch	<i>Carduelis chloris</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
	common redpoll	<i>Carduelis flammea</i>	Introduced and Naturalised	Introduced and Naturalised		P	P	P	P	P	P	P	P	P	P	

Māori Name	Common Name	Scientific Name	National Conservation Status	Regional Conservation Status	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
kōurarini	goldfinch	<i>Carduelis carduelis</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P
hurukōwhai	yellowhammer	<i>Emberiza citrinella</i>	Introduced and Naturalised	Introduced and Naturalised	P	P	P	P	P	P	P	P	P	P	P	P

Appendix Two

This table provides a summary of the annual mean number of birds detected per bird count for each native forest bird species that has been recorded in Wellington City since 2011. Species are ordered from the most to least frequently encountered in 2022. Green rows denote species for which mean encounter rates have increased significantly since 2011; yellow rows denote species with stable encounter rates between 2011 and 2022 and red rows represent those species for which encounter rates have declined since 2011. Section 3.2 of this report provides a more detailed picture of the temporal trends in encounter rates for each individual species on this list.

Species	Average number of birds observed at each station											
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
Tūī	1.35	0.77	2.16	1.74	2.60	2.36	2.07	2.49	2.23	2.98	2.35	2.50
Tauhou / silvereye	1.86	2.38	2.03	2.20	2.05	2.40	2.34	2.02	1.98	2.06	1.71	1.69
Riroriro / grey warbler	0.84	1.24	1.29	1.05	1.45	1.06	1.29	1.26	0.87	1.01	1.38	1.18
Pīwakawaka / New Zealand fantail	0.35	0.22	0.23	0.32	0.21	0.28	0.35	0.36	0.43	0.42	0.48	0.52
Kākā	0.10	0.07	0.12	0.10	0.24	0.15	0.22	0.26	0.26	0.35	0.37	0.36
Kererū	0.07	0.03	0.11	0.08	0.08	0.18	0.13	0.13	0.20	0.20	0.24	0.21
Pīpīwharau / shining cuckoo	0.17	0.23	0.24	0.19	0.18	0.20	0.23	0.21	0.06	0.08	0.11	0.18
Tieke / North Island saddleback	0.07	0.02	0.03	0.02	0.06	0.06	0.10	0.09	0.06	0.06	0.04	0.07
Kākāriki / red-crowned parakeet	0.01	0.01	0.01	0.03	0.08	0.07	0.06	0.09	0.04	0.06	0.03	0.06
Korimako / bellbird	0.00	0.02	0.03	0.01	0.00	0.01	0.03	0.01	0.05	0.02	0.02	0.06
Pōpokotea / whitehead	0.06	0.09	0.04	0.06	0.05	0.05	0.08	0.07	0.11	0.05	0.06	0.05

Species	Average number of birds observed at each station											
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
Kōtare / New Zealand kingfisher	0.03	0.04	0.03	0.04	0.04	0.02	0.03	0.06	0.06	0.04	0.05	0.02
Kārearea / New Zealand falcon	0.00	0.01	0.01	0.00	0.00	0.01	0.01	0.01	0.00	0.01	0.03	0.02
Toutouwai / North Island robin	0.01	0.01	0.01	0.00	0.00	0.00	0.00	0.01	0.02	0.00	0.02	0.02
Tītipounamu / rifleman	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Hihi / stitchbird	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00
Miromiro / tomtit	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Koekoeā / long-tailed cuckoo	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Ruru / morepork	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00